

Free Local Delivery | 24 Hour Turnaround Time | Competitive Price Match

Podiatry Compounding

Wound Care

- Phenytoin 5 % / Misoprostol 0.0024 %
Topical Gel
- Phenytoin 2 % / Misoprostol 0.0024 % /
Lidocaine 2 % / Bupivacaine HCl 0.2 % /
Diphenhydramine HCl 1 % / Aloe Vera 0.2 %
Polyox Bandage
- Ketoprofen 2 % / Lidocaine 2 % / Misoprostol
0.0024 % / Phenytoin 2 % / Aloe Vera 0.2 %
Topical Cream

Diabetic Toes

- Nifedipine 4 % Topical Lipoderm
- Pentoxifylline (5 %, 10 %, 15 %) Lipoderm
Transdermal
- Ketamine 10 % / Gabapentin 6 % / Clonidine
0.2 % / Nifedipine 2 % Topical Lipoderm

Raynaud's Phenomenon

- Nifedipine 16 % Topical Lipoderm (Toe Butter)
- Nifedipine 4 % Topical Lipoderm
- Pentoxifylline (5 %, 10 %, 15 %) Lipoderm
Transdermal

Rough/Dry Feet

- Urea 25 % Topical Cream
- Urea 8 % Topical Cream

Amputation Pain

- Ketoprofen 20 % / Ibuprofen 2 % / Ketamine
2 % / Clonidine 0.2 % / Amitriptyline HCl 2 %
/ Guaifenesin 2 % Topical Lipoderm
- Baclofen 5 % / Ketoprofen 10 % / Lidocaine 5 %
/ Gabapentin 5 % Topical Lipoderm
- Ketamine HCl 5 % / Gabapentin 10 % / Clonidine
HCl 0.2 % / Baclofen 2 % Topical Lipoderm
- Ketoprofen 5 % / Cyclobenzaprine HCl 0.5 % /
Lidocaine HCl 5 % / Bupivacaine HCl 1 %
Topical Lipoderm

Inflammation from Injury

- Piroxicam 5 % in Lipoderm Transdermal Gel
- Piroxicam 2 % Topical Anhydrous Lipoderm
- Diclofenac Sodium 10 % in Lipoderm
Transdermal Gel
- Ketoprofen 10 % Topical Lipoderm
- Ibuprofen 20 % in Lipoderm Transdermal Gel

**FIRST-TIME PATIENTS:
\$10 OFF***

*mention this ad